

## **Syngenta**

Report generated on: December 4, 2023

Visit our data catalog at: <a href="https://microdata.fao.org/index.php">https://microdata.fao.org/index.php</a>

### Identification

SURVEY ID NUMBER
EGY\_2014-2019\_GGP-P\_v01\_EN\_M\_A\_OCS

TITI F

Good Growth Plan, 2014-2019

#### **COUNTRY**

Name	Country code
Egypt	EGY

#### STUDY TYPE

Agricultural Survey [ag/oth]

#### **ABSTRACT**

Syngenta is committed to increasing crop productivity and to using limited resources such as land, water and inputs more efficiently. Since 2014, Syngenta has been measuring trends in agricultural input efficiency on a global network of real farms. The Good Growth Plan dataset shows aggregated productivity and resource efficiency indicators by harvest year. The data has been collected from more than 4,000 farms and covers more than 20 different crops in 46 countries. The data (except USA data and for Barley in UK, Germany, Poland, Czech Republic, France and Spain) was collected, consolidated and reported by Kynetec (previously Market Probe), an independent market research agency. It can be used as benchmarks for crop yield and input efficiency.

KIND OF DATA

Sample survey data [ssd]

**UNIT OF ANALYSIS** 

Agricultural holdings

### Scope

#### **NOTES**

Data was collected on the usage of inputs, such as crop protection products, chemical fertilizer, seeding rates, labor hours, machinery usage hours, and marketable crop yield on a per hectare basis.

#### **TOPICS**

Topic	Vocabulary
Agriculture & Rural Development	FAO
Environment	FAO
Agricultural input efficiency	FAO

### **KEYWORDS**

Keyword
Input efficiency
Crop productivity
Agriculture
The Good Growth Plan

### Coverage

GEOGRAPHIC COVERAGE

National Coverage

## **Producers and sponsors**

#### PRIMARY INVESTIGATORS

Name	
Syngenta	

#### **PRODUCERS**

Name	Role
Kynetec	Technical assistance

## Sampling

#### SAMPLING PROCEDURE

#### A. Sample design

Farms are grouped in clusters, which represent a crop grown in an area with homogenous agro- ecological conditions and include comparable types of farms. The sample includes reference and benchmark farms. The reference farms were selected by Syngenta and the benchmark farms were randomly selected by Kynetec within the same cluster.

#### B. Sample size

Sample sizes for each cluster are determined with the aim to measure statistically significant increases in crop efficiency over time. This is done by Kynetec based on target productivity increases and assumptions regarding the variability of farm metrics in each cluster. The smaller the expected increase, the larger the sample size needed to measure significant differences over time. Variability within clusters is assumed based on public research and expert opinion. In addition, growers are also grouped in clusters as a means of keeping variances under control, as well as distinguishing between growers in terms of crop size, region and technological level. A minimum sample size of 20 interviews per cluster is needed. The minimum number of reference farms is 5 of 20. The optimal number of reference farms is 10 of 20 (balanced sample).

### C. Selection procedure

The respondents were picked randomly using a "quota based random sampling" procedure. Growers were first randomly selected and then checked if they complied with the quotas for crops, region, farm size etc. To avoid clustering high number of interviews at one sampling point, interviewers were instructed to do a maximum of 5 interviews in one village.

BF Screened from Egypt were from Assuit, Menia, Banisuif, Sharqiya, Ismaliaand were selected based on the following criterion:

- BACKGROUND: Open field tomatoes
- Flood irrigation
- Ploughing with a tractor or manually (e.g. with a hoe)
- Usage of chemical and/or organic fertilizers
- Selling the harvest is the main after harvest activity

### **Data Collection**

#### DATES OF DATA COLLECTION

Start	End
2014	2019

## **Questionnaires**

#### **QUESTIONNAIRES**

Data collection tool for 2019 covered the following information:

#### (A) PRE- HARVEST INFORMATION

PART I: Screening

PART II: Contact Information PART III: Farm Characteristics a. Biodiversity conservation

- b. Soil conservation
- c. Soil erosion
- d. Description of growing area
- e. Training on crop cultivation and safety measures

PART IV: Farming Practices - Before Harvest

- a. Planting and fruit development Field crops
- b. Planting and fruit development Tree crops
- c. Planting and fruit development Sugarcane
- d. Planting and fruit development Cauliflower
- e. Seed treatment

#### (B) HARVEST INFORMATION

PART V: Farming Practices - After Harvest

- a. Fertilizer usage
- b. Crop protection products
- c. Harvest timing & quality per crop Field crops
- d. Harvest timing & quality per crop Tree crops
- e. Harvest timing & quality per crop Sugarcane
- f. Harvest timing & quality per crop Banana
- g. After harvest

PART VI - Other inputs - After Harvest

- a. Input costs
- b. Abiotic stress
- c. Irrigation

See all questionnaires in external materials tab

# **Data Processing**

### DATA EDITING

Data processing:

Kynetec uses SPSS (Statistical Package for the Social Sciences) for data entry, cleaning, analysis, and reporting. After collection, the farm data is entered into a local database, reviewed, and quality-checked by the local Kynetec agency. In the case of missing values or inconsistencies, farmers are re-contacted. In some cases, grower data is verified with local experts (e.g. retailers) to ensure data accuracy and validity. After country-level cleaning, the farm-level data is submitted to the global Kynetec headquarters for processing. In the case of missing values or inconsistences, the local Kynetec office was recontacted to clarify and solve issues.

#### B. Quality assurance

Various consistency checks and internal controls are implemented throughout the entire data collection and reporting process in order to ensure unbiased, high quality data.

Screening: Each grower is screened and selected by Kynetec based on cluster-specific criteria to ensure a comparable

group of growers within each cluster. This helps keeping variability low.

- Evaluation of the questionnaire: The questionnaire aligns with the global objective of the project and is adapted to the local context (e.g. interviewers and growers should understand what is asked). Each year the questionnaire is evaluated based on several criteria, and updated where needed.
- Briefing of interviewers: Each year, local interviewers familiar with the local context of farming -are thoroughly briefed to fully comprehend the questionnaire to obtain unbiased, accurate answers from respondents.
- Cross-validation of the answers:
- o Kynetec captures all growers' responses through a digital data-entry tool. Various logical and consistency checks are automated in this tool (e.g. total crop size in hectares cannot be larger than farm size)
- o Kynetec cross validates the answers of the growers in three different ways:
- 1. Within the grower (check if growers respond consistently during the interview)
- 2. Across years (check if growers respond consistently throughout the years)
- 3. Within cluster (compare a grower's responses with those of others in the group)
- o All the above mentioned inconsistencies are followed up by contacting the growers and asking them to verify their answers. The data is updated after verification. All updates are tracked.
- Check and discuss evolutions and patterns: Global evolutions are calculated, discussed and reviewed on a monthly basis jointly by Kynetec and Syngenta.
- Sensitivity analysis: sensitivity analysis is conducted to evaluate the global results in terms of outliers, retention rates and overall statistical robustness. The results of the sensitivity analysis are discussed jointly by Kynetec and Syngenta.
- It is recommended that users interested in using the administrative level 1 variable in the location dataset use this variable with care and crosscheck it with the postal code variable.

## **Data Appraisal**

### DATA APPRAISAL

Due to the above mentioned checks, irregularities in fertilizer usage data were discovered which had to be corrected:

For data collection wave 2014, respondents were asked to give a total estimate of the fertilizer NPK-rates that were applied in the fields. From 2015 onwards, the questionnaire was redesigned to be more precise and obtain data by individual fertilizer products. The new method of measuring fertilizer inputs leads to more accurate results, but also makes a year-on-year comparison difficult. After evaluating several solutions to this problems, 2014 fertilizer usage (NPK input) was reestimated by calculating a weighted average of fertilizer usage in the following years.

# **Access policy**

### **CONTACTS**

Name	Affiliation	Email	URL
The Good Growth Plan team	Syngenta	goodgrowthplan.data@syngenta.com	<u>Link</u>

#### CONFIDENTIALITY

The users shall not take any action with the purpose of identifying any individual entity (i.e. person, household, enterprise, etc.) in the micro dataset(s). If such a disclosure is made inadvertently, no use will be made of the information, and it will be reported immediately to FAO

### ACCESS CONDITIONS

Micro datasets disseminated by FAO shall only be allowed for research and statistical purposes. Users requesting access to any datasets must agree to the following minimal conditions:

- The micro dataset will only be used for statistical and/or research purposes;
- Any results derived from the micro dataset will be used solely for reporting aggregated information, and not for any specific

individual entities or data subjects;

- The users shall not take any action with the purpose of identifying any individual entity (i.e. person, household, enterprise, etc.) in the micro dataset(s). If such a disclosure is made inadvertently, no use will be made of the information, and it will be reported immediately to FAO;
- The micro dataset cannot be re-disseminated by users or shared with anyone other than the individuals that are granted access to the micro dataset by FAO.

#### CITATION REQUIREMENTS

The Good Growth Plan Progress Data - Productivity 2019. Dataset downloaded from https://microdata.fao.org.

## **Disclaimer and copyrights**

#### DISCLAIMER

The user of the data acknowledges that the original collector of the data, the authorized distributor of the data, and the relevant funding agency bear no responsibility for use of the data or for interpretations or inferences based upon such uses

## **Metadata production**

#### DDI DOCUMENT ID

EGY\_2014-2019\_GGP-P\_v01\_EN\_M\_A\_OCS\_v01

#### **PRODUCERS**

Name	Abbreviation	Affiliation	Role
Office of Chief Statistician	ocs	Food and Agriculture Organization	Metadata producer

#### **DDI DOCUMENT VERSION**

DDI\_EGY\_2014-2019\_GGP-P\_v01\_EN\_M\_A\_OCS\_FAO

# **Data Dictionary**

Data file	Cases	Variables
crop_protection_country_EGY	2008	32
Farm_level_data_EGY	599	32
Fertilizers_EGY	1292	17
global_farm_data_country_EGY	599	295
Location_EGY	699	18
Q382_data_EGY	5723	9
seed_treatment_EGY	295	24